

Data Assimilation and Model Simulations in the California Current

Robert L. Haney
Department of Meteorology (MR/Hy)
Naval Postgraduate School
589 Dyer Rd., Room 254
Monterey, CA 93943-5114
Phone: (831) 656-4053
Fax: (831) 656-3061
e-mail: haneyrl@met.nps.navy.mil
Award #: N0001498WR30093

LONG-TERM GOALS

The long-term goal of my research is to develop, verify, and apply numerical ocean prediction models to eastern boundary coastal regions in order to improve our scientific understanding of the structure and dynamics of such regions.

OBJECTIVES

The broad objective of this research is to aid in the development of a reliable modeling capability for eastern boundary current regions. My specific objective is to carry out and extensively verify several DieCAST model simulations of the annual cycle in the California Current. I also wish to evaluate and apply digital filter initialization (DFI) as a diagnostic tool in numerical ocean prediction.

APPROACH

I intend to carry out increasingly complex numerical simulations of the California Current using the DieCAST regional model. I will then verify the simulations against ONR's extensive eastern boundary current (EBC) data sets. I also intend to evaluate iterative versions of digital filter initialization (DFI) as a diagnostic tool for ocean analysis, and then apply the method to several quasi-synoptic hydrographic data sets from the California Current and the Alboran Sea. Results from the California Current will be compared with those obtained using the iterated geostrophic method (IGM, Allen 1993) and the results from the Alboran Sea will be verified using direct measurements of the vertical velocity determined by quasi-isobaric RAFOS floats.

WORK COMPLETED

I have adapted the DieCAST A-grid, low-dispersion (4th order) regional model (Dietrich 1997) to the California coastal domain and completed two six-year long numerical simulations. Lateral boundary conditions are prescribed from monthly climatological data and from a global ocean model simulation. One of the simulations uses a climatological annual cycle of wind stress forcing, while the other simulation includes an additional, idealized wind stress enhancement near each coastal headland representative of observations (e.g., Enriquez and Friehe 1995; Rogers et al. 1998). Preliminary results

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Data Assimilation and Model Simulations in the California Current				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Meteorology, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

have been published in an NPS Masters Thesis (Donato 1998), while a complete analysis of the simulations is still in progress.

I have applied iterative forms of DFI to several idealized test cases and have determined an optimal version of the method for use in diagnostic studies. DFI is now being applied to data sets from the California Current (ONR's CTZ and EBC programs) and the Alboran Sea (EC's OMEGA project).

I contributed to two studies of the Alboran Sea that were published this year, and I collaborated in a study of stochastic resonance in the thermohaline circulation that we recently submitted for publication.

RESULTS

Using climatological forcing, our DieCAST model simulation reproduces many of the main features of the observed annual cycle of the California Current including the strengthening of the coastal jet in spring and the weakening of the jet in autumn and winter. Coastal eddies in the simulation form primarily off the major headlands, especially Cape Mendocino and Point Arena. As a result, a region of maximum eddy kinetic energy (EKE), originally formed in the upper ocean over the continental slope in late spring, migrates westward on a seasonal time scale. At the same time, the EKE spreads vertically into the deep ocean, decreasing the EKE west of about 126W. This result identifies a non-dissipative process that can account for the pronounced decrease of surface EKE west of 126W recently documented in the literature (Kelly et al. 1998, Strub and James 1998). The additional forcing by the headland wind jets is found to produce both local and remote changes to the simulated annual cycle.

The optimal form of DFI was shown to be able to diagnose the 3-dimensional circulation and vertical velocity in an idealized growing baroclinic frontal wave with a relative accuracy greater than 90%. At the same time, it is shown that accurate measurements of the currents are needed in order to diagnose the mesoscale circulation on space scales as short as the Rossby radius. Using DFI, a closed cyclonic eddy observed off Point Arena California was diagnosed to exhibit relatively small vertical velocities on the scale of the eddy itself, but to have very significant vertical velocities ($w \sim 30 \text{ m/d}$ at 100 m) associated with sub-eddy scale meanders in the otherwise circular flow around the eddy.

In the Alboran Sea studies, we used remotely sensed and in situ data to document a new mode of synoptic scale variability in the Western Alboran Gyre and to interpret the variability dynamically. In our thermohaline circulation study we showed that a small periodic variation in the fresh water fluxes, as small as 5% of the mean fresh water flux, in the presence of background variability (noise), can induce quasi-periodic transitions between two stable states of the thermohaline circulation.

IMPACT/APPLICATIONS

Our California Current annual cycle simulations are among the most advanced ones to date. Because of the excellent observational data sets being used, the model verifications will represent the standards against which other models will be evaluated in the future.

TRANSITIONS

Our research on data assimilation and simulations in the California Current is in broad support of the efforts at FNMOC (M. Clancy) and NRL-Stennis (J. Kindle) to develop a real-time ocean analysis and forecasting capability, including biology, for this region. This support manifests itself in close coordination, and timely information exchange on such topics as model properties, experimental set-up, forcing data, regional modeling difficulties, comparison and exchange of results, methods of model verification, and so forth.

The DFI methodology has been accepted as the major diagnostic tool for estimating the ageostrophic circulation and vertical motion in the EC-sponsored OMEGA project. This multinational European project completed six quasi-synoptic oceanographic surveys in the western and central Mediterranean last year.

RELATED PROJECTS

I am collaborating with J. Barth (OSU) in his application of the DFI diagnostic method to the EBC synoptic survey data.

I am collaborating with D. Dietrich (MSU-CAST) in a DieCAST model hindcast of the Santa Barbara Channel driven by high resolution wind stress data from COAMPS. I am also providing the COAMPS wind stress data to Leo Oey for a similar SBC hindcast using the Princeton Ocean Model (POM).

I am collaborating with A. Viudez and J. Tintore (UIB, Spain) in diagnostic studies of mesoscale variability in the Alboran Sea and modeling studies of climatic variability of the thermohaline circulation.

REFERENCES

Allen, J. S., 1993: Iterated geostrophic intermediate models. *J. Phys. Oceanogr.*, **23**, 2447-2461.

Dietrich, D. E., 1997: Application of a modified Arakawa "A" grid ocean model having reduced numerical dispersion to the Gulf of Mexico circulation. *Dyn. Atmos. and Oceans*, **27**, 201-217.

Donato, J., 1998: The seasonal cycle of mesoscale variability in the California Current: A DieCAST model simulation. M. S. Thesis, Department of Meteorology, Naval Postgraduate School, Monterey, CA, Sep. 1998, 68pp.

Enriquez, A. G., and C. A. Friehe, 1995: Effects of wind stress and wind stress curl variability on coastal upwelling. *J. Phys. Oceanogr.*, **25**, 1651-1671.

Kelly, K. A., R. C. Beardsley, R. Limeburner, K. H. Brink, J. D. Paduan and T. K. Chereskin, 1998: Variability of the near-surface eddy kinetic energy in the California Current based on altimetric, drifter and moored current data. *J. Geophys. Res.*, **103**, 13067-13083.

Rogers, D. P., C. E. Dorman, K. A. Edwards, I. M. Brooks, W. K. Melville, S. D. Burk, W. T. Thompson, T. Holt, L. M. Strom, M. Tjernstrom, B. Grisogono, J. M. Bane, W. A. Nuss, B. M. Morley and A. J. Schanot, 1998: Highlights of Coastal Waves 1996. *Bull. Am. Meteor. Soc.*, **79**, 1307-1326.

Strub, P. T. and C. James, 1998: Altimeter-derived variability of surface velocities in the California Current system: 2. Seasonal circulation and eddy statistics. *Deep Sea Res.* (in press).

PUBLICATIONS

Donato, J., 1998: The seasonal cycle of mesoscale variability in the California Current: A DieCAST model simulation. M. S. Thesis, Department of Meteorology, Naval Postgraduate School, Monterey, CA, Sep. 1998, 68pp.

Viudez, A., R. L. Haney and J. Vazquez-Cuervo, 1998: The deflection and division of an oceanic baroclinic jet by a coastal boundary: A case study in the Alboran Sea. *J. Phys. Oceanogr.*, **28**, 289-303.

Viudez, A., J.-M. Pinot and R. L. Haney, 1998: On the upper layer circulation in the Alboran Sea. *J. Geophys. Res.*, **103**, 21653-21666.

Viudez, A., R. L. Haney and J. T. Allen, 1998: A study of the balance of horizontal momentum in a vertical shearing current. *J. Phys. Oceanogr.* (submitted).

Shearman, R. K., J. A. Barth, J. S. Allen and R. L. Haney, 1998: Diagnosis of the three-dimensional circulation in mesoscale features with large Rossby number. *J. Phys. Oceanogr.* (submitted).

Velez, P., A. Alvarez, P. Colet, J. Tintore and R. L. Haney, 1998: Stochastic resonance in the global ocean conveyor belt. *Climate Dynamics* (submitted).

Haney, R. L., 1997: Global warming? Invited presentation, Annual Climate Diagnostics and Prediction Workshop, E. O. Lawrence Berkeley National Laboratory, Berkeley. CA, 6-10 Oct., 1997.

Haney, R. L., 1998: The general circulation in the California Current: Observations and simulations. Invited presentation, Akio Arakawa Fest: Symposium on General Circulation Model Development: Past, Present, and Future, UCLA, 20-22 Jan., 1998.

Haney, R. L., D. E. Dietrich and R. A. Hale, 1998: The role of headland wind jets in the annual cycle of the California Current system. Paper presented at the 1998 Ocean Sciences Meeting, San Diego, CA, 9-13 Feb., 1998.